



Single-stage and two-stage total failure-based group-sampling plans for the Weibull distribution under neutrosophic statistics

Muhammad Aslam¹ · G. Srinivasa Rao² · Nasrullah Khan³

Received: 13 July 2020 / Accepted: 4 December 2020 / Published online: 5 January 2021
© The Author(s) 2021

Abstract

If the sample or population has vague, inaccurate, unidentified, deficient, indecisive, or fuzzy data, then the available sampling plans could not be suitable to use for decision-making. In this article, an improved group-sampling plan based on time truncated life tests for Weibull distribution under neutrosophic statistics (NS) has been developed. We developed improved single and double group-sampling plans based on the NS. The proposed design neutrosophic plan parameters are obtained by satisfying both producer's and consumer's risks simultaneously under neutrosophic optimization solution. Tables are constructed for the selected shape parameter of Weibull distribution and various combinations of neutrosophic group size. The efficiency of the proposed group-sampling plan under the neutrosophic statistical interval method is also compared with the crisp method grouped sampling plan under classical statistics.

Keywords Neutrosophic non-linear problem · Neutrosophic statistics · Consumer's risk · Producer's risk · Group acceptance sampling · Average sample number

List of symbols

T_{Ni}	Neutrosophic random variable
T_L	Determinate parts of nrv
$T_U I_N$	Indeterminate parts of nrv
b_N	Neutrosophic shape parameter
σ_N	Neutrosophic scale parameter
μ_N	Neutrosophic mean life
t_{0N}	Neutrosophic termination time
μ_{0N}	Neutrosophic target mean life
n_N	Neutrosophic random sample
g_N	Neutrosophic groups
c_N	Neutrosophic acceptance number
$L(p)$	Lot acceptance probability

Introduction

The sampling plans are important instruments to judge the quality of manufactured products and components in many fields, including the packing industry, food industry, electrical engineering, aeronautical engineering, and automobile engineering. For instance, before final delivery of items to the customers, it is essential to confirm whether the product meets the specification laid down by the company. The decision of selecting reliable and higher quality products can be made through a statistical sampling approach. In the method of sampling, if the aim is to decide whether to accept or reject many manufactured products, this type of assessment system is usually called acceptance sampling; for more details, see Montgomery [36]. Thus, an acceptance-sampling plan is an investigative method in statistical quality control to make a judgment on submitted manufactured product lots, either to accept or reject. If the quality of the product is the lifetime of the product, then a sample of such products considered exemplify of the observed lifetimes of the goods are put further for testing. When a decision to accept or reject the lot, subject to the risks associated with the two types of errors (rejecting a good lot/accepting a bad lot), is possible, then such a procedure is known as acceptance-sampling plans based on life test. It needs specifications of a certain probability model prevailing the life of the products.

✉ Muhammad Aslam
aslam_ravian@hotmail.com

G. Srinivasa Rao
gaddesrao@gmail.com

Nasrullah Khan
nasrullah.khan@uvas.edu.pk

¹ Department of Statistics, Faculty of Science, King Abdulaziz University, Jeddah 21551, Saudi Arabia

² Department of Mathematics and Statistics, The University of Dodoma, P.O. Box: 259, Dodoma, Tanzania

³ Department of Statistics, University of Veterinary and Animal Sciences (Jhang Campus), Lahore, Pakistan

Acceptance sampling was initiated by Dodge and Romig [22], and this method played a significant role in the manufacturing industry as well as quality management in business development. Bray and Lyon [19] have given the application of acceptance-sampling plans in the food industry. Broadly, acceptance-sampling plans are classified as variables sampling plans and attribute sampling plans. Variables sampling plans are useful when quality characteristics are measurable. When quality characteristics are not measurable, they can be classified as conforming or nonconforming.

A chief advantage in an acceptance-sampling plan is to optimize both the time and cost necessary for the conclusion about the acceptance or rejection of the submitted lot of products in quality control or reliability tests. The acceptance-sampling plans could also give the preferred safeguard to both producers and consumers. Thus, in decision-making using acceptance-sampling schemes, both producer's and consumer's risks are required. An acceptance-sampling plan, which gives protection to both producer's and consumer's risks, is known as a well-designed plan. In life testing experiments, the sample size directly influences the cost of experimentation. Hence, a sampling plan is said to be more economical if it gives the smaller sample size, and to meet these situations, researchers are using more frequently single sampling plans. Whereas, if it is not possible to get the decision-based on the first sample, a double sampling plan can be employed. A generalized sampling plan is known as a double sampling plan and it performs better than a single sampling plan with respect to the sample size. Hence, there is a need to developing an edition of a double sampling plan for life test using groups, which will be called a two-stage group sampling in this paper. As compared to the single sampling plan, the two-stage sampling plan is more complex to handle. Some of the references on single and double acceptance-sampling plans under various life distributions based on truncated life tests can be seen; for example, Epstein [23], Goode and Kao [24], Gupta and Groll [26], Tsai and Wu [51], Balakrishnan et al. [15], Kantam et al. [33], Baklizi [13], Baklizi and El Masri [14], Aslam [2], Aslam and Jun [8], Aslam et al. [9, 11].

The main goal of acceptance-sampling plans in quality control is to minimize the sample size to save cost, time, and efforts. To achieve this, sometimes, an experimenter can test multiple items in practice, because testing time and cost can be saved by testing those items simultaneously. The items in a tester can be regarded as a group and the number of items in a group is called the group size. An acceptance-sampling plan based on such groups of items is called a group acceptance-sampling plan (GASP). If the GASP is used in conjunction with truncated life tests, it is called a GASP based on truncated life test, assuming that the lifetime of the product follows a certain probability distribution. For such a type of test, the determination of the sample size is equivalent to

determine the number of groups. Pascual and Meeker [37] and Jun et al. [31] initiated these group-sampling plans. Subsequently, many authors concentrated on grouped sampling plan for the truncated life test; see example, Aslam et al. [10], Aslam and Jun [7], Rao [43, 44], Aslam et al. [11]. Rao and Rameshnaidu [45], Aslam et al. [12] proposed a new type of grouped sampling plan for truncated life tests, which are based on single and double group-sampling plans under the total number of failures from all groups under testing.

The group-sampling plans developed by the aforementioned researchers under classical statistics can only be applied when there is no uncertainty in the sample or parameters. If the observations or parameters are uncertain or indeterminate, the more popular approach is based on the fuzzy method. Hence, the sampling plans designed using fuzzy logic can be applied to make a decision on a lot of the product. More details about the fuzzy approach in different areas (such as engineering, project management, pattern recognition, transmission systems, multiobjective optimization, etc.) can be seen in Beg and Tabasam [16], Biswas et al. [18], Huang and Wei [27], Majumdar and Samanta [35], Peng and Dai [39], Peng and Liu [40], Peng and Yang [41], Peng [38], and Zhang and Xu [57]. In recent years, more attention has been given to developing a sampling plan using the fuzzy approach [1, 20, 21, 28–30, 32, 34, 46, 50, 52–54, 56].

The neutrosophic logic was introduced by Smarandache [47]. The neutrosophic logic is the extension of the fuzzy logic, and is a combination of measure of truth, a measure of falsehood, and measure of indeterminacy. The fuzzy logic is unable to give information about the measure of indeterminacy. Gulistan and Salma [25] discussed the application of neutrosophic logic using complex fuzzy sets. More information about neutrosophic logic can be seen in Yang [55], Peng and Dai [42], Zhang et al. [58], and Zhang et al. [59].

In recent years, neutrosophic logic is used as the generalization of fuzzy logic. Based on the neutrosophic logic, Smarandache [48, 49] introduced the neutrosophic statistics (NS) as an extension of classical statistics. The classical statistics cannot be applied when the data are obtained from the complex process and have indeterminate or uncertain values. In such a case, to analyze the data, neutrosophic statistics can be applied. The neutrosophic statistics provide information about the measure of indeterminacy which classical statistics does not provide. Therefore, neutrosophic statistics can be considered as the generalization of classical statistics. The neutrosophic statistics approach in sampling schemes was extensively used by different researchers including Aslam and Arif [5], Aslam and Raza [6], and Aslam [4]. Mostly, in earlier literature, Aslam et al. [12] studied a traditional improved group-sampling plans for Weibull distribution, while there is no research on improved group-sampling plans for Weibull distribution under the neutrosophic statistics. Aslam and Arif [5] worked on the sampling plan using the



idea of sudden death testing. Aslam [4] proposed the attribute sampling plan using neutrosophic statistics. Aslam et al. [3] proposed the group-sampling plan under neutrosophic statistics.

In Aslam et al. [3] plan, the number of failures from each group is recorded and a lot of the product is rejected if in any group the number of failures is larger than the allowed number of failures. Note here that the plan proposed by Aslam et al. [3] can be improved by making the decision about the lot based on the total number of failures from all groups. In the literature, the group-sampling plan under neutrosophic statistics based on the total number of failures is not available. To address this gap, this paper proposes a single and a double group-sampling plan using the neutrosophic statistics. In the proposed sampling plans, the decision about the acceptance or rejection of the lot is made on the basis of the total number of failures from all groups. We assume that the lifetime of the manufactured goods follows the neutrosophic Weibull distribution. The proposed plan will be more efficient and effective than the crisp method of competitive sampling plans in terms of the average sample number. The rest of the paper is organized as follows: “[Design of single group-sampling plan for Weibull distribution under neutrosophic statistics](#)” deals with a single group-sampling plan for Weibull distribution based on neutrosophic statistics and compares it with the classical single group-sampling plan. In “[Comparison with classical single group-sampling plan](#)”, a design of a two-stage group-sampling plan for Weibull distribution under neutrosophic statistics is developed and compared with the classical two-stage group-sampling plan. Finally, in “[Design of two-stage group-sampling plan for Weibull distribution under neutrosophic statistics](#)”, some closing remarks are given.

Design of single group-sampling plan for Weibull distribution under neutrosophic statistics

Assume that the lifetime of the neutrosophic random variable $T_N \in [T_L, T_U]$ where T_L denotes the determinate part and T_U denotes the indeterminate part follows the neutrosophic Weibull distribution. The neutrosophic random variable (nrv) in neutrosophic form can be written as follows: $T_N = T_L + T_U I_N; I_N \in [I_L, I_U]$, where T_L and $T_U I_N$ are determinate and indeterminate parts of nrv, respectively. Note here that $I_N \in [I_L, I_U]$ represents the indeterminate interval. The nrv reduces to the random variable under classical statistic if no indeterminate observation is in the data. The neutrosophic Weibull distribution was developed by Aslam and Arif [5] and neutrosophic cumulative distribution function (ncdf) is given below:

$$F_N(t_N; b_N, \sigma_N) = 1 - \exp\left(-\left(t_N/\sigma_N\right)^{b_N}\right); t_N \geq 0, b_N \in [b_L, b_U], \sigma_N \in [\sigma_L, \sigma_U], \quad (1)$$

where $b_N \in [b_L, b_U]$ is the neutrosophic shape parameter and $\sigma_N \in [\sigma_L, \sigma_U]$ is the neutrosophic scale parameter. The mean lifetime σ_N of a neutrosophic Weibull distribution is $\mu_N = (\sigma_N/b_N)\Gamma(1/b_N)$. The neutrosophic form of ncdf and mean under indeterminacy can be written as, respectively, follows:

$$F_N(t_N; b_N, \sigma_N) = F(t; b, \sigma) + b_N I_{NF}; I_N \in [I_{LF}, I_{UF}]. \quad (2)$$

$$\mu_N = \mu + c_N I_{N\mu}; I_{N\mu} \in [I_{L\mu}, I_{U\mu}]. \quad (3)$$

Note here that $F(t; b, \sigma)$ and μ denote the cumulative distribution function (cdf) and mean under classical statistics, respectively. Also, $b_N I_{NF}; I_N \in [I_{LF}, I_{UF}]$ and $a_N I_{N\mu}; I_N \in [I_{L\mu}, I_{U\mu}]$ denote the corresponding indeterminate parts.

The unknown neutrosophic scale parameter can be expressed in terms of the mean lifetime of a neutrosophic Weibull distribution as $\sigma_N = \mu_N(b_N/\Gamma(1/b_N))$. Let t_{0N} is denoted as specified experiment neutrosophic termination time, and μ_{0N} is denoted as a neutrosophic target mean life. Thus, it is convenient to express specified experiment neutrosophic termination time as a multiple of the neutrosophic target mean life. That is, we consider $t_{0N} = a\mu_{0N}$, where a is an experiment neutrosophic termination ratio. Hence, the failure probability of an item before the experiment time t_{0N} is obtained by putting $t_{0N} = a\mu_{0N}$ and $\mu_N = (\sigma_N/b_N)\Gamma(1/b_N)$ into Eq. (1) can be expressed as follows:

$$p_N = 1 - \exp\left(-a^{b_N}(\mu_{0N}/\mu_N)^{b_N}(\Gamma(1/b_N)/b_N)^{b_N}\right). \quad (4)$$

The probability of failure of an item in neutrosophic form can be written as:

$$p_N = p + d_N I_{Np}; I_{Np} \in [I_{Lp}, I_{Up}]. \quad (5)$$

Note here that p denotes the probability of failure of an item under classical statistics and $d_N I_{Np}; I_{Np} \in [I_{Lp}, I_{Up}]$ represents its indeterminate part.

The systematic procedure for a single group-sampling plan for Weibull distribution using neutrosophic statistics is as follows:

- Step 1. Select a neutrosophic random sample of size n_N from a lot.
- Step 2. Allot r_N items to each of g_N groups or testers, such that $n_N = r_N g_N$.
- Step 3. Conduct the test for r_N items before the termination time t_{0N} .

Step 4. The lot will be accepted if the total number of failures from g_N groups is smaller than or equal to c_N before termination time t_{0N} , otherwise reject the lot.

The proposed neutrosophic single group-sampling plan becomes a classical single group-sampling plan if $r_N = r$, $g_N = g$, and $c_N = c$. In this plan, we are interested to find the design parameters like number of neutrosophic groups g_N and the neutrosophic acceptance number c_N , in addition, to satisfying both producer's risk (α) and consumer's risk (β) for specified values of the group sizes and true quality levels. Based on Step 4, the lot acceptance probability for the proposed plan is given by:

$$L(p_N) = \sum_{i=0}^{c_N} \binom{r_N g_N}{i} p_N^i (1 - p_N)^{r_N g_N - i}, \quad (6)$$

where p_N is given in Eq. (4).

The lot acceptance probability in the form of the indeterminate interval can be given as:

$$L(p_N) = L(p) + e_N I_{NL}; I_{NL} \in [I_{LL}, I_{UL}], \quad (7)$$

where $L(p)$ denote the lot acceptance probability under classical statistics and $e_N I_{NL}; I_{NL} \in [I_{LL}, I_{UL}]$ shows the indeterminate part. Equation (7) reduces to lot acceptance probability under classical statistics when $I_L = 0$. The plan parameters are obtained using a two-point approach where producers aim that the probability of acceptance should be greater than $1 - \alpha$ at the acceptable quality limit (AQL), say p_{1N} and consumer's wish that the probability of acceptance should be smaller than β at the limiting quality level (LQL), say p_{2N} . Therefore, we obtain the design neutrosophic parameters by solving the following two inequalities:

$$L(p_{1N}) = \sum_{i=0}^{c_N} \binom{r_N g_N}{i} p_{1N}^i (1 - p_{1N})^{r_N g_N - i} \geq 1 - \alpha \quad (8)$$

$$L(p_{2N}) = \sum_{i=0}^{c_N} \binom{r_N g_N}{i} p_{2N}^i (1 - p_{2N})^{r_N g_N - i} \leq \beta. \quad (9)$$

If k_1 is the AQL as mean ratio at the producer's risk and k_2 is the LQL at the consumer's risk, then design neutrosophic parameters can be found by satisfying the following inequalities:

$$L(p_{1N} | \mu_N / \mu_{0N} = k_1) \geq 1 - \alpha. \quad (10)$$

$$L(p_{2N} | \mu_N / \mu_{0N} = k_2) \leq \beta. \quad (11)$$

The above-mentioned inequalities are used to find the plan parameters of the proposed plan using the grid search method. Grid search is a process that searches exhaustively

through a manually specified subset of the hyperparameter space of the targeted algorithm for more details refer to Bergstra and Bengio [17]. In grid search, several combinations are found that meet the given conditions, we choose those values of parameters where n or average sample number (ASN) is minimum. During the simulation, it is noted that several combinations of the parameters exist which satisfied the given conditions. The plan parameters where $g_N \in [g_L, g_U]$ is minimum. Here, we regard as $k_2 = 1$. Tables 1, 2 were built for neutrosophic Weibull distribution with neutrosophic shape parameter $b_N \in [1.9, 2.1]$ at $\beta \in [0.25, 0.10, 0.05, 0.01]$, mean ratios $k_1 \in [2, 4, 6, 8, 10]$, two group sizes $r_N \in [4, 6]$ and $r_N \in [10, 12]$, and two termination times $a = 0.5, 1.0$. From tables, we noticed the following tendency:

- i. When the consumer's risk value decreases, the number of groups (g_N) is increased.
- ii. When the mean ratio value increases, the number of groups (g_N) is decreased.
- iii. The number of groups (g_N) decreases with the increase of r_N . That is, e.g., when other parametric values are fixed at $\beta = 0.25$, $\mu_N / \mu_{0N} = 4$, $b_N \in [1.9, 2.1]$ from Tables 1 for $r_N \in [4, 6]$ with group size $g_N \in [5, 7]$, whereas from Tables 2 for $r_N \in [10, 12]$ with group size $g_N = [2, 4]$.

Comparison with classical single group-sampling plan

In this subsection, a comparison is made between the crisp method classical single group-sampling plan proposed by Aslam et al. [12] and a single group-sampling plan under neutrosophic statistics. The proposed sampling plan is more economical than the crisp method sampling plan due to less number of groups. The comparison between the proposed sampling plan and the crisp method grouped sampling plans is given in Table 3. From Table 3, it is clear that the proposed group-sampling plan significantly decreases the number of groups required as compared with the crisp method group-sampling plan. In particular, when the mean ratio is small, the proposed plan shows better performance than the crisp method plan, whereas if the mean ratio is large, the proposed plan is reduced to the crisp method plan. For example, if we consider at $\beta = 0.25$, $\alpha = 0.05$, $b = 2$, $r = 4$, $a = 1.0$, and $r_I = 4$, the crisp method plan proposed by Aslam et al. [12] needs $g = 2$ and $c = 2$, whereas in the proposed plan under neutrosophic statistics needs a number of groups and acceptance numbers in indeterminate form as: $g_N = 1 + 3I_N$, $I_N \in [0, 0.6]$, and $c_N = 1 + 5I_N$; $I_N \in [0, 0.8]$. Here, we note that the chance of indeterminacy in the selection of group size and acceptance numbers is 60% and 80%, respectively. We note that the



Table 1 Proposed single group sampling plan for Weibull distribution under neutrosophic statistics $r_N \in [4, 6]$, $b_N \in [1.9, 2.1]$, and various $I_N \in [I_L, I_U]$

β	$\mu_N/\mu_{0N} = k_1$	$a = 0.5$			$a = 1.0$		
		g_N	c_N	$L(p_{1N})$	g_N	c_N	$L(p_{1N})$
0.25	2	[5, 7]	[2, 3]	[0.903, 0.906]	[-, -]	[-, -]	[-, -]
	4	[4, 6]	[1, 2]	[0.976, 0.995]	[1, 3]	[1, 5]	[0.983, 1]
	6	[4, 6]	[1, 2]	[0.994, 1]	[1, 3]	[1, 5]	[0.996, 1]
	8	[4, 6]	[1, 3]	[0.998, 1]	[1, 3]	[1, 3]	[0.999, 1]
	10	[4, 6]	[1, 2]	[0.999, 1]	[1, 3]	[1, 5]	[0.999, 1]
0.10	2	[10, 12]	[4, 5]	[0.931, 0.923]	[-, -]	[-, -]	[-, -]
	4	[5, 7]	[1, 2]	[0.963, 0.992]	[2, 4]	[1, 3]	[0.931, 0.984]
	6	[5, 7]	[1, 2]	[0.991, 0.999]	[2, 4]	[1, 2]	[0.983, 0.991]
	8	[5, 7]	[1, 2]	[0.997, 1]	[2, 4]	[1, 6]	[0.994, 1]
	10	[5, 7]	[1, 2]	[0.999, 1]	[2, 4]	[2, 3]	[1, 1]
0.05	2	[-, -]	[-, -]	[-, -]	[-, -]	[-, -]	[-, -]
	4	[6, 8]	[1, 3]	[0.949, 0.999]	[2, 4]	[1, 6]	[0.931, 1]
	6	[6, 8]	[1, 3]	[0.988, 1]	[2, 4]	[1, 2]	[0.983, 0.991]
	8	[6, 8]	[1, 2]	[0.996, 1]	[2, 4]	[1, 3]	[0.994, 1]
	10	[6, 8]	[1, 2]	[0.998, 1]	[2, 4]	[1, 3]	[0.997, 1]
0.01	2	[-, -]	[-, -]	[-, -]	[-, -]	[-, -]	[-, -]
	4	[8, 10]	[1, 2]	[0.915, 0.979]	[3, 5]	[2, 4]	[0.974, 0.993]
	6	[8, 10]	[1, 3]	[0.978, 1]	[3, 5]	[2, 6]	[0.997, 1]
	8	[8, 10]	[1, 2]	[0.992, 1]	[3, 5]	[2, 5]	[0.999, 1]
	10	[8, 10]	[1, 3]	[0.997, 1]	[3, 5]	[1, 4]	[0.994, 1]

[-, -] shows that parameters do not exist

Table 2 Proposed single group-sampling plan for Weibull distribution under neutrosophic statistics $r_N \in [10, 12]$, $b_N \in [1.9, 2.1]$, and various $I_N \in [I_L, I_U]$

β	$\mu_N/\mu_{0N} = k_1$	$a = 0.5$			$a = 1.0$		
		g_N	c_N	$L(p_{1N})$	g_N	c_N	$L(p_{1N})$
0.25	2	[2, 4]	[2, 5]	[0.903, 0.986]	[2, 4]	[6, 11]	[0.927, 0.912]
	4	[2, 4]	[2, 4]	[0.997, 1]	[1, 3]	[2, 11]	[0.985, 1]
	6	[2, 4]	[2, 4]	[1, 1]	[1, 3]	[3, 12]	[1, 1]
	8	[2, 4]	[2, 4]	[1, 1]	[1, 3]	[1, 10]	[0.99, 1]
	10	[2, 4]	[1, 4]	[0.999, 1]	[1, 3]	[2, 8]	[1, 1]
0.10	2	[4, 6]	[4, 7]	[0.931, 0.99]	[2, 4]	[6, 12]	[0.927, 0.956]
	4	[2, 4]	[1, 2]	[0.963, 0.988]	[1, 3]	[2, 11]	[0.985, 1]
	6	[2, 4]	[1, 2]	[0.991, 0.999]	[1, 3]	[1, 5]	[0.973, 1]
	8	[2, 4]	[1, 4]	[0.997, 1]	[1, 3]	[1, 9]	[0.99, 1]
	10	[2, 4]	[1, 2]	[0.999, 1]	[1, 3]	[2, 3]	[1, 1]
0.05	2	[6, 8]	[6, 9]	[0.952, 0.994]	[2, 4]	[6, 12]	[0.927, 0.956]
	4	[3, 5]	[1, 2]	[0.924, 0.979]	[1, 3]	[2, 9]	[0.985, 1]
	6	[3, 5]	[1, 2]	[0.981, 0.998]	[1, 3]	[2, 7]	[0.998, 1]
	8	[3, 5]	[1, 2]	[0.993, 1]	[1, 3]	[2, 6]	[1, 1]
	10	[3, 5]	[1, 4]	[0.997, 1]	[1, 3]	[2, 12]	[1, 1]
0.01	2	[8, 10]	[7, 10]	[0.923, 0.989]	[-, -]	[-, -]	[-, -]
	4	[5, 7]	[2, 3]	[0.959, 0.99]	[2, 4]	[4, 5]	[0.996, 0.986]
	6	[4, 6]	[1, 3]	[0.967, 1]	[1, 3]	[1, 9]	[0.973, 1]
	8	[4, 6]	[1, 4]	[0.988, 1]	[1, 3]	[1, 10]	[0.99, 1]
	10	[4, 6]	[1, 3]	[0.995, 1]	[1, 3]	[1, 4]	[0.996, 1]

[-, -] shows parameters do not exist

values of g and c from the crisp method sampling plan lie in the group size indeterminacy interval [1, 3] and acceptance

number indeterminacy interval [1, 5]. From this comparison, it can be seen that the new algorithm calculates more

Table 3 Comparison of group-sampling plan under classical statistics and under neutrosophic statistics when $r_N \in [4, 6]$ and $b_N \in [1.9, 2.1]$ with classical for $r=4$, $b=1.9$

β	$\mu_N/\mu_{0N} = k_1$	Classical method with $a = 1.0$			Neutrosophic method with $a = 1.0$		
		g	c	$L(p_1)$	g_N	c_N	$L(p_{1N})$
0.25	2	4	6	0.9783	[–, –]	[–, –]	[–, –]
	4	2	2	0.9922	[1, 3]	[1, 5]	[0.983, 1]
	6	2	2	0.9991	[1, 3]	[1, 5]	[0.996, 1]
	8	2	2	0.9998	[1, 3]	[1, 3]	[0.999, 1]
	10	2	1	0.9973	[1, 3]	[1, 5]	[0.999, 1]
0.10	2	5	7	0.9742	[–, –]	[–, –]	[–, –]
	4	2	2	0.9922	[2, 4]	[1, 3]	[0.931, 0.984]
	6	2	2	0.9991	[2, 4]	[1, 2]	[0.983, 0.991]
	8	2	1	0.9939	[2, 4]	[1, 6]	[0.994, 1]
	10	2	1	0.9973	[2, 4]	[2, 3]	[1, 1]
0.05	2	6	8	0.9714	[–, –]	[–, –]	[–, –]
	4	3	2	0.9741	[2, 4]	[1, 6]	[0.931, 1]
	6	2	1	0.9828	[2, 4]	[1, 2]	[0.983, 0.991]
	8	2	1	0.9939	[2, 4]	[1, 3]	[0.994, 1]
	10	2	1	0.9973	[2, 4]	[1, 3]	[0.997, 1]
0.01	2	8	10	0.9684	[–, –]	[–, –]	[–, –]
	4	3	2	0.9741	[3, 5]	[2, 4]	[0.974, 0.993]
	6	3	2	0.9967	[3, 5]	[2, 6]	[0.997, 1]
	8	3	2	0.9993	[3, 5]	[2, 5]	[0.999, 1]
	10	3	2	0.9998	[3, 5]	[1, 4]	[0.994, 1]

[–, –] shows that parameters do not exist

compactly a set of parameters, containing the values of the classical one. In addition, the proposed sampling plan provides the smaller values of g_N as compared to the crisp method plan.

Design of two-stage group-sampling plan for Weibull distribution under neutrosophic statistics

We present the design of a two-stage group-sampling plan for Weibull distribution under neutrosophic statistics in this section. Aslam et al. [12] studied similar plans for classical statistics and we used the same algorithm under neutrosophic statistics.

The systematic procedure to obtain design parameters in this plan is given below:

First stage:

- Step 1. Select at random the first sample of size n_{1N} from a lot.
- Step 2. Assign r items to each of g_{1N} groups (or testers), such that $n_{1N} = rg_{1N}$.
- Step 3. Conduct the test for r items before the termination time t_{0N} .

Step 4. The lot will be accepted if the total number of failures from g_{1N} groups is smaller than or equal to c_{1aN} .

Step 5. Terminate the test and reject the lot when the total number of failures is greater than or equal to c_{1rN} ($> c_{1aN}$) before termination time t_{0N} . Otherwise, go to the second stage.

Second stage:

- Step 1. Choose the second random sample of size n_{2N} from a lot.
- Step 2. Assign r items to each of g_{2N} groups, such that $n_{2N} = rg_{2N}$.
- Step 3. Conduct the test for r items before the termination time t_{0N} .
- Step 4. The lot will be accepted if the total number of failures from g_{1N} and g_{2N} groups is smaller than or equal to c_{2aN} ($\geq c_{1aN}$) before termination time t_{0N} ; otherwise, reject the lot.

The projected two-stage group-sampling plan for Weibull distribution under neutrosophic statistics is differentiated by five design parameters, namely g_{1N} , g_{2N} , c_{1aN} , c_{1rN} , and c_{2aN} . The proposed plan can be reduced to a single group-sampling plan based on neutrosophic statistics outlined in “[Design of single group-sampling plan for Weibull distribution under](#)

neutrosophic statistics” when $c_{1rN} = c_{1aN} + 1$. As a result, the total number of failures from g_{1N} groups (denoted by X_{1N}) follows a binomial distribution with parameters n_{1N} and p_N . Hence, the probabilities of acceptance and rejection of the lot at the first stage under the proposed two-stage group-sampling plan are given below:

$$P_{aN}^{(1)} = \sum_{i=0}^{c_{1aN}} \binom{r g_{1N}}{i} p_N^i (1 - p_N)^{r g_{1N} - i}. \quad (12)$$

$$P_{rN}^{(1)} = \sum_{i=c_{1rN}}^{n_{1N}} \binom{r g_{1N}}{i} p_N^i (1 - p_N)^{r g_{1N} - i}. \quad (13)$$

Also, the probability of acceptance of lot from the second stage if the decision has not been completed at the first stage and the total number of failures from g_{1N} , g_{2N} groups (denoted by X_{2N}) is smaller than or equal to c_{2aN} . Therefore:

$$\begin{aligned} P_{aN}^{(2)} &= P(c_{1aN} + 1 \leq X_{1N} \leq c_{1rN} - 1, X_{1N} + X_{2N} < c_{2aN}) \\ &= \sum_{x=c_{1aN}+1}^{c_{1rN}-1} \binom{r g_{1N}}{x} p_N^x (1 - p_N)^{r g_{1N} - x} \\ &\quad \left[\sum_{i=0}^{c_{2aN}-x} \binom{r g_{2N}}{i} p_N^i (1 - p_N)^{r g_{2N} - i} \right]. \end{aligned} \quad (14)$$

Hence, the probability of lot acceptance for the proposed two-stage group-sampling plan based on neutrosophic statistics is given below:

$$L(p_N) = P_{aN}^{(1)} + P_{aN}^{(2)}.$$

The lot acceptance probability in the form of the indeterminate interval can be given as:

$$L(p_N) = L(p) + f_N I_N; I_N \in [I_L, I_U], \quad (15)$$

where $L(p)$ denotes the lot acceptance probability for Aslam et al.'s [11] plan and $f_N I_N$, $I_N \in [I_L, I_U]$ denotes the indeterminate part. The optimum plan parameters are obtained by minimizing the average sample number (ASN), Aslam et al. [11]. Hence, the optimum plan parameters are a solution to the following neutrosophic non-linear optimization problem:

$$\text{Minimize } \text{ASN}(p_{2N}) = r_N g_{1N} + r_N g_{2N} (1 - P_{aN}^{(1)} - P_{rN}^{(1)}). \quad (16)$$

Subject to

$$L(p_{1N}) = P_{aN}^{(1)}(p_{1N}) + P_{aN}^{(2)}(p_{2N}) \geq 1 - \alpha. \quad (17)$$

$$L(p_{2N}) = P_{aN}^{(1)}(p_{1N}) + P_{aN}^{(2)}(p_{2N}) \leq \beta. \quad (18)$$

The above-mentioned inequalities are used to find the plan parameters of the proposed plan using a grid search method. During the simulation, it is noted that several combinations of the parameters exist, which satisfies the given conditions. The plan parameters which have minimum ASN is selected and reported. The design parameters are displayed in Tables 4, 5 for neutrosophic Weibull distribution with neutrosophic shape parameter $b_N \in [1.9, 2.1]$ at $\beta = [0.25, 0.10, 0.05, 0.01]$, mean ratios $k_1 = [2, 4, 6, 8, 10]$, two group sizes $r = 10, 20$, and two termination times, $a = 0.5$ and 1.0 . Smarandache [49] suggested that de-neutrosophication can be done using the average of each interval or minimum/maximum value of intervals can be considered. From tables, we noticed the following tendency by considering the minimum values of intervals:

- The ASN value decreases as k_1 increases from 2 to 6 for the fixed value of β .
- The ASN value decreases as a increases from 0.5 to 1.0 for the fixed value of b_N .
- The values of ASN increases as the values of r increase from 10 to 20 when other parameters are fixed.

Comparison with classical two-stage group-sampling plan

In this subsection, a comparison is made between a single group-sampling plan under neutrosophic statistics, the crisp method, classical two-group-sampling plan proposed by Aslam et al. [12], and proposed two group-sampling plans under neutrosophic Statistics. The proposed sampling plan is said to be more effective than the crisp method sampling plan if it gives less ASN and, hence, the plan is more economical. For comparison, we considered common parameters $\beta = 0.25$, $\alpha = 0.05$, $b = 2$, $a = 0.5$, and $r_I = 4$ for three plans. From Tables 2 and 4, we noticed that the neutrosophic interval is smaller in the two-group-sampling plan under neutrosophic statistics than in a single group-sampling plan under neutrosophic statistics. For example from Table 4 when the neutrosophic shape parameter $b_N \in [1.9, 2.1]$, the

Table 4 Proposed two-stage group-sampling plan for Weibull distribution under neutrosophic

β	$\mu_N/\mu_{0N} = k_1$	$a = 0.5$						$a = 1.0$					
		c_{1aN}	c_{2aN}	c_{1rN}	g_{1N}	g_{2N}	ASN	c_{1aN}	c_{2aN}	c_{1rN}	g_{1N}	g_{2N}	ASN
0.25	2	[1, 4]	[3, 6]	[6, 8]	[2, 4]	[1, 2]	[23.064, 40.474]	[3, 5]	[5, 7]	[4, 9]	[1, 2]	[1, 2]	[10.797, 21.966]
	4	[0, 4]	[3, 6]	[6, 8]	[2, 4]	[1, 2]	[22.64, 40.001]	[1, 5]	[4, 7]	[6, 9]	[1, 2]	[1, 2]	[11.034, 20.002]
	6	[0, 4]	[2, 6]	[5, 8]	[2, 4]	[1, 2]	[21.323, 40]	[0, 5]	[2, 7]	[4, 9]	[1, 2]	[1, 2]	[12.326, 20]
	8	[0, 4]	[2, 6]	[5, 8]	[2, 4]	[1, 2]	[20.789, 40]	[1, 5]	[4, 7]	[5, 9]	[1, 2]	[1, 2]	[10.096, 20]
	10	[1, 4]	[3, 6]	[4, 9]	[2, 4]	[1, 2]	[20.013, 40]	[0, 5]	[2, 7]	[7, 9]	[1, 2]	[1, 2]	[10.955, 20]
0.10	2	[0, 4]	[4, 6]	[6, 8]	[3, 5]	[1, 2]	[38.192, 51.091]	[-, -]	[-, -]	[-, -]	[-, -]	[-, -]	[-, -]
	4	[0, 3]	[2, 5]	[4, 7]	[2, 4]	[1, 3]	[22.64, 40.019]	[2, 5]	[5, 7]	[4, 9]	[1, 2]	[1, 2]	[10.153, 20.002]
	6	[0, 3]	[2, 5]	[4, 7]	[2, 4]	[1, 3]	[21.323, 40.001]	[1, 5]	[3, 7]	[6, 9]	[1, 2]	[1, 2]	[10.267, 20]
	8	[0, 3]	[2, 5]	[5, 7]	[2, 4]	[1, 3]	[20.789, 40]	[0, 5]	[3, 7]	[7, 9]	[1, 2]	[1, 2]	[11.421, 20]
	10	[0, 3]	[2, 5]	[5, 7]	[2, 4]	[1, 3]	[20.523, 40]	[1, 5]	[3, 7]	[7, 9]	[1, 2]	[1, 2]	[10.043, 20]
0.05	2	[3, 5]	[5, 7]	[4, 9]	[5, 7]	[1, 2]	[51.584, 71.37]	[-, -]	[-, -]	[-, -]	[-, -]	[-, -]	[-, -]
	4	[0, 3]	[3, 5]	[4, 7]	[3, 5]	[1, 2]	[33.686, 50.03]	[1, 5]	[4, 7]	[6, 9]	[1, 2]	[1, 2]	[11.034, 20.002]
	6	[1, 3]	[3, 5]	[5, 7]	[3, 5]	[1, 2]	[30.191, 50.001]	[0, 5]	[2, 7]	[5, 9]	[1, 2]	[1, 2]	[12.326, 20]
	8	[1, 3]	[3, 5]	[5, 7]	[3, 5]	[1, 2]	[30.068, 50]	[1, 5]	[4, 7]	[6, 9]	[1, 2]	[1, 2]	[10.096, 20]
	10	[0, 3]	[2, 5]	[4, 7]	[3, 5]	[1, 2]	[30.775, 50]	[1, 5]	[4, 7]	[4, 9]	[1, 2]	[1, 2]	[10.043, 20]
0.01	2	[3, 3]	[5, 7]	[4, 9]	[6, 8]	[1, 2]	[61.893, 88.402]	[-, -]	[-, -]	[-, -]	[-, -]	[-, -]	[-, -]
	4	[0, 2]	[2, 4]	[4, 6]	[3, 5]	[2, 4]	[37.371, 50.522]	[0, 5]	[4, 7]	[6, 9]	[1, 2]	[1, 2]	[14.357, 20.002]
	6	[0, 2]	[2, 4]	[4, 6]	[3, 5]	[2, 4]	[33.834, 50.05]	[0, 5]	[4, 7]	[5, 9]	[1, 2]	[1, 2]	[12.326, 20]
	8	[0, 2]	[2, 4]	[4, 6]	[3, 5]	[2, 4]	[32.319, 50.009]	[0, 5]	[2, 7]	[4, 9]	[1, 2]	[1, 2]	[11.421, 20]
	10	[0, 2]	[2, 4]	[4, 6]	[3, 5]	[2, 4]	[31.55, 50.002]	[0, 5]	[4, 7]	[4, 9]	[1, 2]	[1, 2]	[10.955, 20]

statistics $r=10$, $b_N=[1.9, 2.1]$ and various $I_N \in [I_L, I_U]$

[-, -] shows that parameters do not exist

neutrosophic ASN in a two-stage group-sampling plan under neutrosophic statistics gives [23.064, 40.474], whereas from Table 2 in single group, sampling plan under neutrosophic statistics gives $[2 \times 10 = 20, 4 \times 12 = 48]$. Furthermore, the two-stage group-sampling plan under neutrosophic statistics shows better performance than the crisp method classical two-stage group-sampling plan proposed by Aslam et al. [12]. For example, in Table 4, when $\mu_N/\mu_{0N} = 2$, $\beta = 0.25$, neutrosophic ASN in a two-stage group-sampling plan under neutrosophic statistics gives [23.064, 40.474], whereas in the double group-sampling plan proposed by Aslam et al. [12] needs an ASN value of 39.8. When $\mu_N/\mu_{0N} = 2$, $\beta = 0.05$, neutrosophic ASN in a two-stage group-sampling plan under neutrosophic statistics gives [51.584, 71.37], whereas in the double group-sampling plan proposed by Aslam et al. [12] needs ASN value of 52.4. When $\mu_N/\mu_{0N} = 2$, $\beta = 0.01$, neutrosophic ASN in a two-stage group-sampling plan under neutrosophic statistics gives [61.893, 88.402], whereas in the double group-sampling plan proposed by Aslam et al. [12] needs ASN value 62.9. From this study, it is clear that the proposed plan has smaller values of ASN as compared to the crisp method plan proposed by Aslam et al. [12]. Hence, the proposed neutrosophic approach is more cost-effective and saving of experiment time.

Conclusions

In this article, two types of group acceptance-sampling plans using neutrosophic statistics are considered for Weibull distribution. The neutrosophic plan parameters are developed for both single and two-stage group-sampling plans using neutrosophic statistics. Various tables are provided for industrial applications. A comparative study is also carried out, and it shows that the proposed two-stage group-sampling plan under neutrosophic statistics performs better than the crisp method classical double group-sampling plan and the proposed single group-sampling plan under neutrosophic statistics based on the ASN values. From the comparison, it is found that the proposed sampling plans have smaller values of group size as compared to the crisp method sampling plans. In addition, the comparisons show that the proposed sampling plans are effective, flexible, and adequate to be applied when indeterminacy is presented. The proposed plan can be applied in the industry when there is uncertainty in observations or parameters or both. The proposed group-sampling plan using repetitive sampling and multiple dependent state sampling can be considered as future research.

Table 5 Proposed two-stage group-sampling plan for Weibull distribution under neutrosophic statistics $r=20$, $b_N=[1.9, 2.1]$, and various $I_N \in [I_L, I_U]$

β	$\mu_N/\mu_{0N} = k_1$	$a = 0.5$						$a = 1.0$					
		c_{1aN}	c_{2aN}	c_{1rN}	g_{1N}	g_{2N}	ASN	c_{1aN}	c_{2aN}	c_{1rN}	g_{1N}	g_{2N}	ASN
0.25	2	[0, 4]	[4, 6]	[5, 8]	[1, 2]	[1, 2]	[33.619, 40.948]	[1, 15]	[11, 17]	[17, 19]	[1, 2]	[1, 2]	[38.39, 40.014]
	4	[0, 4]	[2, 6]	[5, 8]	[1, 2]	[1, 2]	[25.281, 40.002]	[3, 15]	[8, 17]	[10, 19]	[1, 2]	[1, 2]	[20.452, 40]
	6	[0, 4]	[2, 6]	[5, 8]	[1, 2]	[1, 2]	[22.646, 40]	[0, 15]	[2, 17]	[5, 19]	[1, 2]	[1, 2]	[28.223, 40]
	8	[0, 4]	[2, 6]	[5, 8]	[1, 2]	[1, 2]	[21.577, 40]	[1, 13]	[6, 15]	[8, 18]	[1, 2]	[1, 2]	[20.734, 40]
	10	[0, 4]	[2, 6]	[7, 10]	[1, 2]	[1, 2]	[21.047, 40]	[0, 12]	[2, 17]	[12, 19]	[1, 2]	[1, 2]	[23.636, 40]
0.10	2	[0, 3]	[4, 6]	[4, 8]	[1, 2]	[1, 2]	[33.549, 43.277]	[2, 14]	[12, 17]	[13, 19]	[1, 2]	[1, 2]	[35.381, 40.047]
	4	[0, 3]	[2, 5]	[5, 7]	[1, 2]	[1, 2]	[25.281, 40.025]	[2, 15]	[5, 17]	[16, 19]	[1, 2]	[1, 2]	[21.935, 40]
	6	[0, 3]	[2, 5]	[4, 7]	[1, 2]	[1, 2]	[22.646, 40.001]	[4, 15]	[6, 17]	[17, 19]	[1, 2]	[1, 2]	[20.003, 40]
	8	[0, 3]	[3, 5]	[5, 7]	[1, 2]	[1, 2]	[21.577, 40]	[1, 13]	[6, 17]	[10, 19]	[1, 2]	[1, 2]	[20.734, 40]
	10	[0, 3]	[2, 5]	[4, 7]	[1, 2]	[1, 2]	[21.047, 40]	[3, 14]	[5, 17]	[12, 19]	[1, 2]	[1, 2]	[20.001, 40]
0.05	2	[0, 7]	[6, 9]	[5, 11]	[2, 4]	[1, 2]	[57.526, 80.696]	[1, 14]	[14, 16]	[11, 18]	[1, 2]	[1, 2]	[38.388, 40.047]
	4	[0, 2]	[2, 4]	[4, 6]	[1, 2]	[1, 2]	[25.281, 40.283]	[0, 15]	[4, 17]	[17, 19]	[1, 2]	[1, 2]	[33.631, 40]
	6	[0, 2]	[2, 4]	[4, 6]	[1, 2]	[1, 2]	[22.646, 40.026]	[2, 15]	[4, 17]	[8, 19]	[1, 2]	[1, 2]	[20.292, 40]
	8	[0, 2]	[2, 4]	[4, 6]	[1, 2]	[1, 2]	[21.577, 40.005]	[0, 12]	[2, 15]	[15, 17]	[1, 2]	[1, 2]	[25.281, 40]
	10	[0, 2]	[2, 4]	[4, 6]	[1, 2]	[1, 2]	[21.047, 40.001]	[3, 12]	[6, 14]	[4, 16]	[1, 2]	[1, 2]	[20.001, 40]
0.01	2	[4, 7]	[6, 9]	[5, 11]	[3, 5]	[1, 2]	[62.497, 102.175]	[4, 13]	[13, 15]	[7, 18]	[1, 2]	[1, 2]	[26.22, 40.144]
	4	[1, 5]	[4, 7]	[5, 9]	[2, 4]	[1, 2]	[42.473, 80.006]	[0, 13]	[7, 15]	[13, 17]	[1, 2]	[1, 2]	[33.631, 40]
	6	[1, 5]	[4, 7]	[5, 9]	[2, 4]	[1, 2]	[40.653, 80]	[1, 13]	[3, 16]	[11, 18]	[1, 2]	[1, 2]	[21.903, 40]
	8	[1, 5]	[4, 7]	[6, 9]	[2, 4]	[1, 2]	[40.236, 80]	[0, 12]	[2, 14]	[6, 19]	[1, 2]	[1, 2]	[25.281, 40]
	10	[0, 5]	[4, 8]	[11, 15]	[2, 4]	[1, 2]	[42.039, 80]	[0, 11]	[2, 13]	[13, 15]	[1, 2]	[1, 2]	[23.636, 40]

Acknowledgements The authors are deeply thankful to the editor and reviewers for their valuable suggestions to improve the quality of this paper.

Compliance with ethical standards

Conflict of interest None.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Alaeddini A, Ghazanfari M, Nayeri MA (2009) A hybrid fuzzy-statistical clustering approach for estimating the time of changes in fixed and variable sampling control charts. *Inf Sci* 179:1769–1784
- Aslam M (2007) A double acceptance sampling plan based on truncated life testing Rayleigh distribution. *Eur J Sci Res* 17(4):605–611
- Aslam M (2019) A new attribute sampling plan using neutrosophic statistical interval method. *Complex Intell Syst* 5:1–6
- Aslam M, Arif O (2018) Testing of grouped product for the Weibull distribution using neutrosophic statistics. *Symmetry* 10:403
- Aslam M, Jun C-H (2009) A group acceptance sampling plan for truncated life test having Weibull distribution. *J Appl Stat* 39:1021–1027
- Aslam M, Jun HC (2010) A double acceptance sampling plan for generalized log-logistic distribution with known shape parameters. *J Appl Stat* 37(3):405–414
- Aslam M, Raza MA (2018) Design of new sampling plans for multiple manufacturing lines under uncertainty. *Int J Fuzzy Syst* 1–15
- Aslam M, Jun C-H, Ahmad M (2009a) A group sampling plan based on truncated life tests for gamma distributed items. *Pakistan J Stat* 25:333–340
- Aslam M, Jun HC, Ahmad M (2009b) A double acceptance sampling plan based on the truncated life tests in the Weibull model. *J Stat Theo Appl* 8:191–206
- Aslam M, Jun HC, Ahmad M (2010) Design of a time truncated double sampling plan for a general life distribution. *J Appl Stat* 37(8):1369–1379
- Aslam M, Jun CH, Lee H, Ahmad M, Rasool M (2011) Improved group sampling plans based on truncated life tests. *Chil J Stat* 2(1):85–97

12. Aslam M, Jeyadurga P, Balamurali S, Al-Marshadi AH (2019) Time-truncated group plan under the Weibull distribution based on neutrosophic statistics. *Mathematics* 7(10):905
13. Baklizi A (2003) Acceptance sampling plan based on truncated life test in the Pareto distribution of second kind. *Adv Appl Stat* 3(1):33–48
14. Baklizi A, El Masri QEA (2004) Acceptance sampling plan based on truncated life test in the Birnbaum-Saunders model. *Risk Anal* 24(6):1453–1457
15. Balakrishnan N, Leiva V, López J (2007) Acceptance sampling plans from truncated life tests based on the generalized Birnbaum-Saunders distribution. *Commun Stat Simul Comput* 36:643–656
16. Beg I, Tabasam R (2013) TOPSIS for hesitant fuzzy linguistic term sets. *Int J Fuzzy Syst* 28:1162–1171
17. Bergstra J, Bengio Y (2012) Random search for hyper-parameter optimization. *J Mach Learn Res* 13:281–305
18. Biswas P, Pramanik S, Giri BC (2016) TOPSIS method for multi-attribute group decision-making under single-valued neutrosophic environment. *Neural Comput Appl* 27:727–737
19. Bray DF, Lyon DA (1973) Three class attributes plans in acceptance sampling. *Technometrics* 15:575–585
20. Cheng S-R, Hsu B-M, Shu M-H (2007) Fuzzy testing and selecting better processes performance. *Ind Manag Data Syst* 107:862–881
21. Divya P (2012) Quality interval acceptance single sampling plan with fuzzy parameter using Poisson distribution. *Int J Adv Res Technol* 1:115–125
22. Dodge HF, Romig HG (1946) Sampling inspection tables: single and doubling sampling. *J Roy Stat Soc* 109(3):297–298
23. Epstein B (1954) Truncated life tests in the exponential case. *Ann Math Stat* 25(3):555–564
24. Goode HP, Kao JHK (1961) Sampling plans based on the Weibull distribution. In: *Proceeding of the Seventh National Symposium on Reliability and Quality Control*. Philadelphia, pp. 24–40
25. Gulistan M, Salma K (2020) Extensions of neutrosophic cubic sets via complex fuzzy sets with application. *Complex Intell Syst* 6:309–320
26. Gupta SS, Groll PA (1961) Gamma distribution in acceptance sampling based on life tests. *J Am Stat Assoc* 56:942–970
27. Huang YH, Wei GW (2018) TODIM method for Pythagorean 2-tuple linguistic multiple attribute decision making. *J Intell Fuzzy Syst* 35:901–915
28. Jamkhaneh EB, Gildeh BS (2011) Chain sampling plan using Fuzzy probability theory. *J Appl Sci* 11(24):3830–3838
29. Jamkhaneh EB, Gildeh BS (2012) Acceptance double sampling plan using fuzzy Poisson distribution. *World Appl Sci J* 16(11):1578–1588
30. Jamkhaneh EB, Gildeh BS (2013) Sequential sampling plan using fuzzy SPRT. *J Intell Fuzzy Syst* 25:785–791
31. Jun C-H, Balamurali S, Lee S-H (2006) Variables sampling plans for Weibull distributed lifetimes under sudden death testing. *IEEE Trans Reliab* 55:53–58
32. Kahraman C, Bekar ET, Senvar O (2016) A fuzzy design of single and double acceptance sampling plans. *Intelligent decision making in quality management*. Springer, Germany, pp 179–211
33. Kanagawa A, Ohta H (1990) A design for single sampling attribute plan based on fuzzy sets theory. *Fuzzy Sets Syst* 37:173–181
34. Kantam RRL, Rosaiah K, Rao GS (2001) Acceptance sampling based on life tests: log-logistic models. *J Appl Stat* 28:121–128
35. Majumdar P, Samanta SK (2014) On similarity and entropy of neutrosophic sets. *J Intell Fuzzy Syst* 26(3):1245–1252
36. Montgomery DC (2009) *Introduction to statistical quality control*, 6th edn. Wiley, New York
37. Pascual FG, Meeker WQ (1998) The modified sudden death test: planning life tests with a limited number of test positions. *J Test Eval* 26:434–443
38. Peng X (2019) New similarity measure and distance measure for Pythagorean fuzzy set. *Complex Intell Syst* 5:101–111
39. Peng X, Dai J (2018) Approaches to single-valued neutrosophic MADM based on MABAC, TOPSIS and new similarity measure with score function. *Neural Comput Appl* 29(10):939–954
40. Peng XD, Dai JG (2020) A bibliometric analysis of neutrosophic set: two decades review from 1998 to 2017. *Artif Intell Rev* 53:199–255
41. Peng X, Liu C (2017) Algorithms for neutrosophic soft decision making based on EDAS, new similarity measure and level soft set. *J Intell Fuzzy Syst* 32:955–968
42. Peng XD, Yang Y (2015) Information measures for interval valued fuzzy soft sets and their clustering algorithm. *J Comput Appl* 35:2350–2354
43. Rao GS (2009) A group acceptance sampling plans for lifetimes following a generalized exponential distribution. *Econ Qual Control* 24(1):75–85
44. Rao GS (2009) A group acceptance sampling plans based on truncated life tests for Marshall-Olkin extended Lomax distribution. *Electron J Appl Stat Anal* 3(1):18–27
45. Rao GS, Ramesh Naidu C (2015) An exponentiated half logistic distribution to develop a group acceptance sampling plans with truncated time. *J Stat Manage Syst* 18(6):519–531
46. Sadeghpour Gildeh B, Baloui Jamkhaneh E, Yari G (2011) Acceptance single sampling plan with fuzzy parameter. *Iran J Fuzzy Syst* 8:47–55
47. Smarandache F (2014) *Introduction to neutrosophic statistics*. Infinite study, El Segundo, CA, USA
48. Smarandache F (2010) Neutrosophic logic-A generalization of the intuitionistic fuzzy logic. In: *Multispace & Multistructure. Neutrosophic Transdisciplinarity (100 Collected Papers of Science)*; Infinite Study: El Segundo, CA, USA, Vol. 4, p. 396
49. Smarandache F (1998) Neutrosophy. Neutrosophic probability, set, and logic, proquest information & learning. *Ann Arbor Michigan USA* 105:118–123
50. Tamaki F, Kanagawa A, Ohta H (1991) A fuzzy design of sampling inspection plans by attributes. *Jpn J Fuzzy Theo Syst* 3:315–327
51. Tsai TR, Wu SJ (2006) Acceptance sampling based on truncated life tests for generalized Rayleigh distribution. *J Appl Stat* 33:595–600
52. Turanoğlu E, Kaya İ, Kahraman C (2012) Fuzzy acceptance sampling and characteristic curves. *Int J Comput Intell Syst* 5:13–29
53. Uma G, Ramya K (2015) Impact of fuzzy logic on acceptance sampling plans—a Review. *Autom Auton Syst* 7:181–185
54. Venkatesh A, Elango S (2014) Acceptance sampling for the influence of trh using crisp and fuzzy gamma distribution. *Aryabhata J Math Inform* 6:119–124
55. Yang HL, Bao YL, Guo ZL (2018) Generalized interval neutrosophic rough sets and its application in multi-attribute decision-making. *Filomat* 32:11–33
56. Zarandi MF, Alaeddini A, Turksen I (2008) A hybrid fuzzy adaptive sampling—run rules for Shewhart control charts. *Inf Sci* 178:1152–1170
57. Zhang XL, Xu ZS (2014) Extension of TOPSIS to multiple criteria decision making with Pythagorean fuzzy sets. *Int J Fuzzy Syst* 29:1061–1078
58. Zhang C, Li DY, Kang XP, Song D, Sangaiah A, Broumi S (2020) Neutrosophic fusion of rough set theory: an overview. *Comput Ind* 115:103117
59. Zhang C, Li DY, Liang JY (2020) Multi-granularity three-way decisions with adjustable hesitant fuzzy linguistic multigranulation decision-theoretic rough sets over two universes. *Inf Sci* 507:665–683

